

underscored material = new  
[bracketed material] = delete

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

HOUSE MEMORIAL 66

**48TH LEGISLATURE - STATE OF NEW MEXICO - FIRST SESSION, 2007**

INTRODUCED BY

Mimi Stewart

A MEMORIAL

SUPPORTING DEPARTMENT OF HEALTH INITIATIVES TO IMPROVE DELIVERY OF RECOMMENDED PAPANICOLAOU AND HUMAN PAPILLOMAVIRUS TESTING, TO MAINTAIN SURVEILLANCE OF CERVICAL PRECANCERS AND TO FORM A HUMAN PAPILLOMAVIRUS-PAPANICOLAOU ADVISORY PANEL.

WHEREAS, cervical cancer represents approximately three percent of cancers in women and two percent of all cancer deaths in New Mexico, and nearly four hundred women in New Mexico are diagnosed with cervical cancer each year and another twenty-six die from the disease; and

WHEREAS, cervical cancer is among the cancers that are most amenable to prevention or secondary intervention through early detection and screening; and

WHEREAS, two-thirds of the New Mexico women newly diagnosed with invasive cervical cancer are under the age of .168401.1

underscoring material = new  
[bracketed material] = delete

1 fifty-five and have families; and

2 WHEREAS, successful and costly cervical cancer screening  
3 and papanicolaou programs coordinated through longstanding  
4 statewide efforts of the department of health and the Indian  
5 health service have reduced the incidence of invasive cervical  
6 cancer in New Mexico to fewer than one hundred new cases  
7 diagnosed each year; and

8 WHEREAS, the majority of women who are diagnosed with  
9 invasive cervical cancer have a history of not being screened  
10 through a papanicolaou test within the past five years; and

11 WHEREAS, extending cervical cancer screening programs to  
12 women living in New Mexico has resulted in a recent convergence  
13 of the incidence rates for cervical cancer among the three main  
14 ethnic groups living in New Mexico; and

15 WHEREAS, mortality rates remain higher among Hispanic and  
16 Native American women when compared to non-Hispanic white  
17 women, presumably due to a number of complex and sometimes  
18 interrelated disparities, including failure to screen; limited  
19 resources; health care access issues, especially in rural  
20 areas; immigrant status; and other cultural and social  
21 barriers; and

22 WHEREAS, twenty distinct high-risk or carcinogenic types  
23 of genital human papillomaviruses primarily transmitted by  
24 skin-to-skin contact during sexual activity are the cause of  
25 virtually all cervical precancers and cancers; and

.168401.1

1           WHEREAS, genital human papillomavirus infections are  
2 extremely common, with approximately forty percent of women  
3 between the ages of eighteen and forty testing positive for  
4 human papillomavirus; and

5           WHEREAS, in certain women, persistent human papillomavirus  
6 infections with high-risk or carcinogenic human papillomavirus  
7 types are at increased risk for cervical precancer and cancer;  
8 and

9           WHEREAS, specific testing for high-risk human  
10 papillomaviruses has been recommended to improve the early  
11 detection of cervical precancers and cancers:

12           A. in women diagnosed with mild or equivocal  
13 papanicolaou test abnormalities; and

14           B. in addition to routine papanicolaou tests in  
15 women thirty years of age and older; and

16           WHEREAS, extremely effective prophylactic human  
17 papillomavirus vaccines have recently become available but are  
18 expensive, costing approximately three hundred sixty dollars  
19 (\$360) per three-dose series; and

20           WHEREAS, many human papillomavirus types not covered by  
21 these vaccines will still cause cervical cancer and, because  
22 the vaccines demonstrate no significant therapeutic effect in  
23 women already infected with human papillomavirus, it is  
24 imperative to remain vigilant in continuing and improving  
25 cervical cancer screening programs; and

underscored material = new  
[bracketed material] = delete

1           WHEREAS, young girls who are sexually naive and have not  
2 been previously exposed to human papilloma viruses would gain  
3 the greatest benefit from human papilloma virus vaccination;  
4 and

5           WHEREAS, improvements in cervical cancer prevention,  
6 including primary vaccines and secondary human papillomavirus  
7 testing, have great potential to reduce the incidence of  
8 cervical cancer, and any relaxation in papanicolaou test  
9 screening due to a false sense of protection in vaccinated  
10 females can result in an increase in the incidence of cervical  
11 cancer rather than the hoped-for reductions; and

12           WHEREAS, the same disparities affecting papanicolaou  
13 screening programs may determine whether young females are  
14 vaccinated with a three-dose human papillomavirus vaccine,  
15 which requires multiple clinic visits to complete the human  
16 papillomavirus vaccine series; and

17           WHEREAS, New Mexico maintains cervical precancer and  
18 cancer screening surveillance and a statewide immunization  
19 system;

20           NOW, THEREFORE, BE IT RESOLVED BY THE HOUSE OF  
21 REPRESENTATIVES OF THE STATE OF NEW MEXICO that it remain  
22 committed to reducing the incidence of cervical precancers and  
23 cancers in New Mexico by supporting the department of health's  
24 efforts directed to improve statewide delivery of papanicolaou  
25 and human papillomavirus testing and the delivery of human

.168401.1

underscoring material = new  
[bracketed material] = delete

1 papilloma virus vaccines to girls between the ages of nine and  
2 fourteen; and

3 BE IT FURTHER RESOLVED that the New Mexico house of  
4 representatives support the department of health's ongoing  
5 efforts to maintain surveillance of cervical precancers and  
6 that it encourage a collaboration of this program with the New  
7 Mexico immunization program's statewide immunization system;  
8 and

9 BE IT FURTHER RESOLVED that the house of representatives  
10 support the formation of a human papilloma virus-papanicolaou  
11 advisory panel composed of experts in human papilloma virus and  
12 cervical cancer screening, immunization and adolescent and  
13 school-based health and representatives of the New Mexico  
14 immunization program, the New Mexico breast and cervical cancer  
15 early detection program, the public health division of the  
16 department of health, statewide school-based health programs  
17 and key health care advocacy groups within the state; and

18 BE IT FURTHER RESOLVED that the New Mexico house of  
19 representatives support a formal collaboration of the New  
20 Mexico health policy commission with the human papillomavirus-  
21 papanicolaou advisory panel; and

22 BE IT FURTHER RESOLVED that the New Mexico house of  
23 representatives support the creation of a research agenda by  
24 the human papillomavirus-papanicolaou advisory panel and the  
25 New Mexico health policy commission to study and identify

.168401.1

underscored material = new  
~~[bracketed material]~~ = delete

1 cervical cancer disparities and cost-effective delivery of  
2 primary and secondary cervical cancer interventions that will  
3 protect and improve the health of New Mexico women; and

4 BE IT FURTHER RESOLVED that copies of this memorial be  
5 transmitted to the department of health, the New Mexico health  
6 policy commission, the interim legislative health and human  
7 services committee and the governor.

8 - 6 -  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25